



ODOUR ASSESSMENT

DEVELOPMENT: Planning Application for extension to existing poultry unit to accommodate a further 32,000 birds.

LOCATION:

**Cwm Farm
Aberhafesp
Newtown
Powys
SY16 3JD**

Prepared for M & JE Jones, DCE and MN Davies

**Roger Parry & Partners LLP Carmarthen
C/O The Estates Office, 20 Salop Road, Oswestry, Shropshire, SY11 2NU
Tel: 01691655334 Fax: 01691 657798
Email: gail@rogerparry.net**

*Also at: 11 Severn Street, Welshpool, Powys, SY21 7AG
Tel: 01938 554499 Fax: 01938 554462
email: welshpool@rogerparry.net*

*Also at: Hogstow Hall, Minsterley, Shrewsbury, SY5 0HZ
Tel: 01743 791336 Fax: 01743 792770
email: mail@rogerparry.net*

Hazard	Receptor	Pathway	Risk Management	Probability of Exposure	Consequence	What is the overall risk
Odour from the manufacture and selection of feed	Neighbouring dwelling houses within 400m of installation	Air	Measures as described in SGN EPR6.09 (Farming) Odour Management at Intensive Livestock Installations. No on site milling and mixing. Feed specifications prepared by the feed compounder's nutrition specialist.	Unlikely	Odour annoyance	Not significant if managed carefully.
Odour from feed delivery or storage	Neighbouring dwelling houses within 400m of installation	Air	Measures as described in SGN EPR6.09 Feed delivery will be sealed to minimise atmospheric dust. Any feed spillage of feed around bin is immediately swept up. Condition of feed bins is checked frequently so that any leaks or damage can be identified.	Unlikely	Odour annoyance	Not significant.
Odours arising from problems with housing ventilation system, inadequate air movement within house leading to high humidity and wet litter. Inadequate system design, causing poor dispersal of odours	Neighbouring dwelling houses within 400m of installation	Air	Measures as described in SGN EPR6.09 The ventilation system will regularly adjusted according to age and requirements of the flock The ventilation will be designed to efficiently remove moisture from the house.	Unlikely	Odour annoyance	Not significant.
Litter management: odours arising from wet litter (see above) The use of insufficient or poor quality litter.	Neighbouring dwelling houses	Air	Measures as described in SGN EPR6.09 Controls on feed and ventilation (see above) help to maintain litter quality. Additional controls include: insulated walls and ceilings to prevent	Unlikely	Odour annoyance	Not significant.

Spillage of water from drinking systems. Disease outbreaks, leading to wet litter.	within 400m of installation		condensation. Concrete floors to prevent water ingress. Stocking density at optimum level to prevent overcrowding. Use health plan, with specialist veterinary input used as necessary			
Carcase disposal: inadequate storage of carcasses on site.	Neighbouring dwelling houses within 400m of installation	Air	Measures as described in SGN EPR6.09 Carcasses are placed in sealed containers awaiting regular collection by a licensed renderer.	Unlikely	Odour annoyance	Not significant.
House clean out (de littering)	Neighbouring dwelling houses within 400m of installation	Air	Litter is carefully placed into trailers close to the house entrances. Trailers are sheeted before transporting off site. De littering will be avoided at weekends during summer months. No litter is stored on site. Litter will be used on operator controlled land in compliance with a manure management plan and used in the Anaerobic digester.	Likely	Odour annoyance	Not significant if carefully managed.
House clean out (Disinfection and fumigation)	Neighbouring dwelling houses within 400m of installation	Air	Carried out by specialist contractors using DEFRA approved chemicals observing correct dilution rates.	Unlikely	Odour annoyance	Not significant if carefully managed.