

## **ODOUR MANAGEMENT PLAN**

DEVELOPMENT:	PLANNING APPLICATION FOR THE HOUSING OF 32,000 FREE RANGE BIRDS (EGG PRODUCTION)
LOCATION:	Frowen Farm Login Whitland Carmarthenshire SA34 0TP
CLIENT:	Anthony and Fairclough
	Roger Parry & Partners LLP Carmarthen C/O The Estates Office, 20 Salop Road, Oswestry, Shropshire, SY11 2NU Tel: 01691655334 Fax: 01691 657798 Email: <u>gail@rogerparry.net</u>
	Also at:11 Severn Street, Welshpool, Powys, SY21 7AG Tel: 01938 554499 Fax: 01938 554462 email: welshpool@rogerparry.net
	Also at: Hogstow Hall, Minsterley, Shrewsbury, SY5 0HZ Tel: 01743 791336 Fax: 01743 792770 email: <u>mail@rogerparry.net</u>

v1 October 2020

The following tables detail the likely sources of Odour from the Poultry Unit at Frowen Farm.

Actions and measures that will be taken to prevent where possible or minimise odour emissions at Frowen Farm.

The site plan submitted as part of the planning application confirms all material storage areas and potential odour emission sources.

The plan is to be reviewed every year from the permit issue date, prior to any changes (major) being made to any operations, or following a complaint, or changes to any management plan the document is to be dated and signed and the area permitting officer notified.

The Free Range Poultry House is to be based within two poultry houses, where the levels of odour would be at their highest concentration. This then results in a lower risk of odour intensity on those closest receptors to the installation.

ODOUR RELATED ISSUES	POTENTIAL RISKS	ACTIONS TAKEN TO REDUCE THE ODOUR AND ODOUR RISKS ON FARM	COMPLETION DATE
Free-range birds	Odour Levels	Twice daily checks coinciding with bird inspections, normally between 07:00-10:00 hours and 16:00-19:00 hours. Any abnormalities to be recorded and investigated.	As and when required
Manufacture and selection of Feed	Milling and mixing of compound feeds. Use of poor quality and odorous ingredients. Feeds which are unbalanced leading to increased excretion of ammonia.	No on site milling to take place. Feed specifications are prepared by a Nutrition Specialist. Feed is supplied only from UKASTA accredited feed mills, so that only approved raw materials are used.	In place
Feed Delivery and Storage	Spillage of feed during delivery and storage. Creation of dust during feed delivery.	Feed delivery systems are sealed to minimise atmospheric dust. Any spillage of feed around the bin is immediately swept up. The condition of the feed bins is regularly checked and	In place

		routine maintenance undertaken if required. Feed deliveries are monitored to avoid dust and spills.	
Ventilation and Dust	<ul> <li>Inadequate air movement in the house, leading to high humidity and wet litter.</li> <li>Inadequate system design, causing poor dispersal of odours.</li> <li>Extraction fans located close to sensitive receptors</li> <li>Dust</li> </ul>	Use of high velocity roof extraction fans in all houses to aid dispersion, checked prior to the cycle commencement by a qualified electrician who will provide 24 hour breakdown cover. The ventilation system is regularly checked and adjusted to match the age and requirements of the flock. The ventilation system is designed to efficiently remove moisture from the house. Humidity recorded daily and maintained in the range of 55- 65% keeping a balance of dry litter and avoiding dust production. Stock inspections carried out only by trained staff to avoid	In Place

		panicking birds therefore creating dust. Dust levels if present controlled during clean out operations.	
Litter Management	Odours arising from wet litter	Controls on feed and ventilation (see above) help to maintain litter quality. Additional controls include; Use of nipple drinkers with drip cups to minimise spillage. Daily checks of drinker height and pressures to avoid capping. Insulated walls and ceilings to prevent condensation. Concrete falls to prevent ingress of water. Stocking levels at optimum to prevent overcrowding. Use of veterinarian bespoke health plan.	In Place

Carcase Disposal	Inadequate storage of carcasses on site.	Carcasses placed into plastic sealed bags, store in sealed, shaded and vermin proof containers away from sensitive receptors. Frequent collection of carcasses. Daily levels of mortalities recorded with abnormalities investigated.	In Place
House Clean Out	Creation of dust associated with litter removal from houses. Use of odorous products during cleaning.	Litter carefully placed into trailers. Trailers sheeted before leaving fill position. Only DEFRA approved and suitable products used. Chemicals containers triple washed at point of use. Wash water sump levels monitored during washing and emptied as required to prevent overfill.	In Place
Used Litter	Storage of used litter on site. Transport of litter and landspreading.	No storage on site at any time, storage in sealed manure store.	In Place

		Manure removal at the end of the cycle. To be stored in sealed building if manure cannot be applied to the land. All trailers sheeted before leaving fill position. Put in a covered manure store. Spread on land if nutrients are required. Contingency – sold to Anaerobic Digestion Plant/local farmers. Avoidance of double handling. Litter used on operators controlled land and sold to third parties.	
Washing operations including vehicles	Loss of dirty water to land or watercourse	Use of specialist contractors for washing operations. Bespoke terminal hygiene program followed, detailing quantities of water and chemical dilution rates. Key staff monitoring washing operations ensuring effective drainage to dirty water tanks.	In Place

		Dirty water tanks monitored	
		during wash down to maintain	
		freeboard.	
		Vehicle washing at designated	
		wash point.	
		All sediment traps and drained	
		cleaned both before and after	
		washing operations.	
Fugitive Emissions	Leaks to doors, bin pipes, feed	Checks to feed storage and	In Place
Fugilive Emissions			III FIACE
	bins, fuel and chemical	fuel pipes.	
	storage.		
		Fuel oil in approved bunded	
		storage tanks.	
		Chemicals – only small	
		amounts of footdip disinfectant	
		held on site in secondary	
		containment.	
Dirty Water Management	Standing dirty water during the	Working areas around houses	In Place
	• • •		
	production cycle or at clean	are concreted and kept	
	out.	cleaning during production	
		cycle.	
	Application of dirty water to		
	land.	At clean out dirty water from	
		houses together with lightly	
		contaminated yard wash is	
		directed to the storage tanks	
		before removed off site and	
		spread to land.	

Abnormal Operations	Water leak/pipe failure	Water consumption monitored daily ensuring early detection, wet area – blanket covered with top up bedding material to prevent increased odour.	In Place
	Bird Health/Sickness	Veterinarian contacted (24 hour care) Litter covered with fresh top up bedding to minimise increased odour until bird health recovered.	
		Abnormal events documented, dated and signed, appropriate plans reviewed and updated to prevent re occurrence.	
Waste Production/Storage	Odour from production or storage areas.	No storage or production of odorous waste on site. Waste management plan in	In Place
		force detailing types and quantities produced along with disposal routes. Records to be kept on site.	
		Cleaning Chemicals Supplied and used by cleaning contractor not stored on site.	

## ODOUR COMPLAINT FORM FROWEN FARM FREE RANGE POULTRY UNIT

**REFERENCE NUMBER** 

DATE RECEIVED

Name and address of the caller	
Telephone number	
Location of caller to installation	
Time and date of complaint of offending odour	

Has the caller any other comments about the odour?	
about the odour?	
Other comments from caller	
Weather conditions	
Wind direction/speed	
Any previous complaints relating to this odour?	

Any other comments	
Other information	
Potential source of odour that could	
give rise to the complaint	
Operations being carried out at time	
of complaint	
<b>—</b> •• •• ••	
Follow up date and time caller contacted	

Action Taken	
Amendments required to the plan	

Completed By

Dated

Signed